



Andropogon gerardii

Native. Ornamental qualities, color changes from green to bluegreen in summer, to deep red-bronze in fall. Sun, H 5-8' (with inflorescence) x W 2'. Zones 3-9.

https://www.flickr.com/people/35478170@N08 Matt Levin

Bluebunch Wheatgrass

Pseudogeneria spicata

Native. Drought tolerant, cool season grass, goes dormant in late summer. Has a nice upright shape with green leaves and golden seed heads. Sun to part shade. H 18-48" (with inflorescence)

Zones 3-7. https://www.flickr.com/people/35478170@N08 Matt Lavin



Blue Fescue

Festuca glauca

Native. Drought tolerant, cool season grass, goes dormant in late summer. Has a nice upright shape with green leaves and golden seed heads. Sun to part shade. H 18-48" (with inflorescence) Zones 3-7. https://commons.wikimedia.org/wiki/Longwood Gardens







Blue Oat Grass

Helictotrichon sempervirens


Native. Ornamental qualities, color changes from green to blue-green in summer, to deep red-bronze in fall. Sun, H 5-8' (with inflorescence) x W 2'. Zones 3-9.

https://www.flickr.com/people/35478170@N08 Matt Lavin



Bottlebrush Squirreltail

Elymus elymoides





Festuca Idahoensis

Native. Small evergreen clump forming grass. Drought tolerant and deer resistant. Excellent for erosion control. Sun. H 12-24" x W 8-10". Zones 4-8.

Photo: Bryan Olsen 2009 via Wikicommons











Indian Rice Grass

Achnatherum hymenoides

Native. Seeds are edible, fixes nitrogen in the soil. Seed pods can be used in flower arrangements. Sun. 8-30" (with inflorescence) x W 18-24". Zones 3-9.

Photo by: Tonuy Frates

Japanese Blood Grass

Imperata cylindrica

Deer resistant, good for erosion control. Cut to ground in late winter before new growth emerges. Full sun, moist well-drained soil. H 1-2'. Zones 5-9.

Photo by: Jim Hood, 2007





Mace Sedge

Carex grayii

Native to eastern North America. Interesting mace-like flower spikes appear in early summer. These can be dried and used in dried flower arrangements. Medium water, tolerates wet soil. Deer resistant. Can re-seed. Full to part sun. Grows 2-3' tall x 1-2' wide. Zones 4-8.

Photo by: Agnieszka Kwiecień, Nova, 2010





Muhly Grass

Muhlenbergia capillaris

A clump-forming warm season grass, emergse late spring to early summer, blooms late summer. When in bloom the inflorescence has the appearance of a pink cloud floating above the foliage. Full sun to part shade. Native to western central U.S. Drought tolerant and deer resistant. Grows 2-3' tall x 2-3' wide. Zones 5-11.

Photo by John Winder, 2010



Karl Foerster

Calamagrostis acutifolia

Dramatic red bronze seed spikes in early summer that last into winter. Clump forming cool season grass. Drought tolerant. Zones 5-11. Sun. H 4-6' (with inflorescence) x W 2-3'.

https://commons.wikimedia.org/w/index.php?title=User:Satin66Flower&action=edit&redlink=1

Little Bluestem

Schizachyrium scoparium

Native. Blue foliage with occasional pinkish-purple striations, topped by fluffy silvery gold or white seed heads. Turns copper color with frost. Sun (has a tendency to flop in too much shade). H 12-24". Zones 3-10.

Photo: Dan Keck via Wikicommons





1974 50 2024 1974 60 2024 1974 70 2024







Full sun to part shade. Warm season grass, new growth in late spring/early summer. Can be cut down in fall or late winter. Drought tolerant. Size is variable from 2' to 7 or 8'. Zones 4-9 https://commons.wikimedia.org/wiki/User:Kenraiz.

Needle and Thread Grass

Hesperostipa comata

Native. Zones 4-8. Full sun. Grows to 40" tall. Drought tolerant. Not deer resistant. Cool season clump-forming grass. Develops unusual long, twisted seed heads.

https://www.flickr.com/people/35478170@N08



Overdam

Calamagrostis arundinacea

Relative of Karl Foerster with similar growth habit. Instead of solid green foliage, Overdam is variegated with white striping on the outer margins. Cool season grass emerges in early spring.

https://www.flickr.com/people/50697352@N00

Pine Grass

Calamagrostis rubescens

Native. Drought tolerant, clump forming. Host plant for a variety of moths and butterflies. Grows wild in the understory of pine forests. H 24-36".

Photo: Matt Levin, 2016 via wikicommons



Limited quantities of all plants, so inclusion on this list is no guarantee of availability





Prairie Dropseed

Sporobolis heterolepsis

Native.,. Benefit to wildlife, host plant for two rare species of leafhoppers. Seeds are eaten by various bird species. Root mass is a nesting site for native bees. Full sun, drought tolerant, deer resistant. H 2-3' with inflorescence. Zones 3-8. Photo: David J. Stang, 2006 via Wiki commonS





Prairie Junegrass

Koeleria macrantha

Native. Cool season bunchgrass, goes dormant in summer. Blooms yellow in early spring. H 6-24" with inflorescence x W 12-24". Zones 4-10. Photo: Matt Lavin 2018 via wikicommons

Switchgrass

Panicum virgatum

Warm season grass, shows new growth in late spring/early summer. Attractive upright grass with airy seed heads. Full sun to part shade, drought tolerant and deer resistant. H 14-55" with inflorescence. Zones 3-9.

Photo: David Stang, 2006 via Wikicommons











Sweetgrass

Hierochloe odorata

Native. Spreads by underground rhizomes. Has a light vanilla fragrance. Considered sacred by many Indigenous Peoples. Prefers full sun to light shade and moist, but not wet, soil. Grows 10-24" tall with inflorescence. Zones 3-9. https://commons.wikimedia.org/wiki/User:SriMesh

Tufted Hair Grass

Deschampsia cespitosa

Native. Clumping ornamental has fine hair-like seed heads that rise 2-3 feet above the clump. Larval plant food for several butterflies. Part sun, drought tolerant. H 12-36" with inflorescence x W 12-24". Zones 4-8. https://commons.wikimedia.org/wiki/User:Lilly_M





Zebra Grass

miscanthus sinensis 'Little Zebra'

Green foliage with distinctive horizontal gold banding. Deer resistant. Cut down in fall or early spring before new growth emerges. Full sun to part shade. Drought tolerant. H 3-4' x W 2-3'. Zones 4-9. https://www.flickr.com/photos/50697352@N00/ F.D. Richards

